



A service of the National Library of Medicine
and the National Institutes of Health

My NCBI [?] [Sign In] [Register]

All Databases PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Books

Search PubMed [x] for campylobacter jejuni sialyltransferase [Go] [Clear] Save Search

Limits Preview/Index History Clipboard Details

Display Summary [x] Show 20 [x] Sort by [x] Send to [x]

About Entrez

Text Version

All: 10 Review: 1 [x]

Items 1 - 10 of 10

One page.

Entrez PubMed

Overview

Help | FAQ

Tutorials

New/Noteworthy [x]

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation

Matcher

Batch Citation

Matcher

Clinical Queries

Special Queries

LinkOut

My NCBI

Related Resources

Order Documents

NLM Mobile

NLM Catalog

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

☐ 1: Sujatha MS, Balaji PV.

Related Articles, Links



Fold-recognition and comparative modeling of human alpha2,3-sialyltransferases reveal their sequence and structural similarities to CstII from Campylobacter jejuni.

BMC Struct Biol. 2006 Apr 19;6:9.

PMID: 16620397 [PubMed - indexed for MEDLINE]

☐ 2: Wokke JH, van den Berg LH.

Related Articles, Links



A way out of the maze: Campylobacter jejuni gene polymorphisms define Guillain-Barre syndrome.

Neurology. 2005 Nov 8;65(9):1350-1. No abstract available.

PMID: 16275819 [PubMed - indexed for MEDLINE]

☐ 3: Yuki N, Odaka M.

Related Articles, Links



Ganglioside mimicry as a cause of Guillain-Barre syndrome.

Curr Opin Neurol. 2005 Oct;18(5):557-61. Review.

PMID: 16155440 [PubMed - indexed for MEDLINE]

☐ 4: Blixt O, Vasiliu D, Allin K, Jacobsen N, Warnock D, Razi N, Paulson JC, Bernatchez S, Gilbert M, Wakarchuk W.

Related Articles, Links



Chemoenzymatic synthesis of 2-azidoethyl-ganglio-oligosaccharides GD3, GT3, GM2, GD2, GT2, GM1, and GD1a.

Carbohydr Res. 2005 Sep 5;340(12):1963-72.

PMID: 16005859 [PubMed - indexed for MEDLINE]

☐ 5: Goodfellow JA, Bowes T, Sheikh K, Odaka M, Halstead SK, Humphreys PD, Wagner ER, Yuki N, Furukawa K, Furukawa K, Plomp JJ, Willison HJ.

Related Articles, Links



Overexpression of GD1a ganglioside sensitizes motor nerve terminals to anti-GD1a antibody-mediated injury in a model of acute motor axonal neuropathy.

J Neurosci. 2005 Feb 16;25(7):1620-8.

PMID: 15716397 [PubMed - indexed for MEDLINE]

☐ 6: Antoine T, Heyraud A, Bosso C, Samain E.





Related Articles, Links



Highly efficient biosynthesis of the oligosaccharide moiety of the GD3 ganglioside by using metabolically engineered Escherichia coli.

Angew Chem Int Ed Engl. 2005 Feb 18;44(9):1350-2. No abstract available.

PMID: 15674992 [PubMed - indexed for MEDLINE]

- ☐ 7: [Chiu CP, Watts AG, Lairson LL, Gilbert M, Lim D, Wakarchuk WW, Withers SG, Strynadka NC.](#) [Related Articles, Links](#)
 Structural analysis of the sialyltransferase CstII from Campylobacter jejuni in complex with a substrate analog.
Nat Struct Mol Biol. 2004 Feb;11(2):163-70. Epub 2004 Jan 18.
PMID: 14730352 [PubMed - indexed for MEDLINE]
- ☐ 8: [Gilbert M, Brisson JR, Karwaski MF, Michniewicz J, Cunningham AM, Wu Y, Young NM, Wakarchuk WW.](#) [Related Articles, Links](#)
 Biosynthesis of ganglioside mimics in Campylobacter jejuni OH4384. Identification of the glycosyltransferase genes, enzymatic synthesis of model compounds, and characterization of nanomole amounts by 600-mhz (1)h and (13)c NMR analysis.
J Biol Chem. 2000 Feb 11;275(6):3896-906.
PMID: 10660542 [PubMed - indexed for MEDLINE]
- ☐ 9: [Eichler E, Jennings HJ, Gilbert M, Whitfield DM.](#) [Related Articles, Links](#)
 Synthesis of a disialylated hexasaccharide of type VIII group B Streptococcus capsular polysaccharide.
Carbohydr Res. 1999 Jun 30;319(1-4):1-16.
PMID: 10520252 [PubMed - indexed for MEDLINE]
- ☐ 10: [Salloway S, Mermel LA, Seamans M, Aspinall GO, Nam Shin JE, Kurjanczyk LA, Penner JL.](#) [Related Articles, Links](#)
 Miller-Fisher syndrome associated with Campylobacter jejuni bearing lipopolysaccharide molecules that mimic human ganglioside GD3.
Infect Immun. 1996 Aug;64(8):2945-9.
PMID: 8757818 [PubMed - indexed for MEDLINE]

Display Show Sort by Send to

[Write to the Help Desk](#)
[NCBI](#) | [NLM](#) | [NIH](#)
[Department of Health & Human Services](#)
[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

Sep 5 2006 08:10:00

EAST Search History

Ref #	Hits	Search Query	DBs	Default Operator	Plurals	Time Stamp
L1	11	((c or campylobacter) adj jejuni) near6 sialyltransferase	USPAT	OR	OFF	2006/09/06 17:16

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:SSSPTA1623SQS

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

* * * * * Welcome to STN International * * * * *
* *

NEWS 1 Web Page URLs for STN Seminar Schedule - N. America
NEWS 2 "Ask CAS" for self-help around the clock
NEWS 3 FEB 27 New STN AnaVist pricing effective March 1, 2006
NEWS 4 MAY 10 CA/CAPplus enhanced with 1900-1906 U.S. patent
records
NEWS 5 MAY 11 KOREAPAT updates resume
NEWS 6 MAY 19 Derwent World Patents Index to be reloaded and
enhanced
NEWS 7 MAY 30 IPC 8 Rolled-up Core codes added to CA/CAPplus and
USPATFULL/USPAT2
NEWS 8 MAY 30 The F-Term thesaurus is now available in CA/CAPplus
NEWS 9 JUN 02 The first reclassification of IPC codes now
complete in
INPADOC
NEWS 10 JUN 26 TULSA/TULSA2 reloaded and enhanced with new search
and
and display fields
NEWS 11 JUN 28 Price changes in full-text patent databases EPFULL
and PCTFULL
NEWS 12 JUL 11 CHEMSAFE reloaded and enhanced
NEWS 13 JUL 14 FSTA enhanced with Japanese patents
NEWS 14 JUL 19 Coverage of Research Disclosure reinstated in DWPI
NEWS 15 AUG 09 INSPEC enhanced with 1898-1968 archive
NEWS 16 AUG 28 ADISCTI Reloaded and Enhanced
NEWS 17 AUG 30 CA(SM)/CAPplus(SM) Austrian patent law changes

NEWS EXPRESS JUNE 30 CURRENT WINDOWS VERSION IS V8.01b, CURRENT
MACINTOSH VERSION IS V6.0c(ENG) AND V6.0Jc(JP),
AND CURRENT DISCOVER FILE IS DATED 26 JUNE 2006.

NEWS HOURS STN Operating Hours Plus Help Desk Availability
NEWS LOGIN Welcome Banner and News Items
NEWS IPC8 For general information regarding STN implementation
of IPC 8

NEWS X25 X.25 communication option no longer available

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

* * * * * STN Columbus * * * * *

FILE 'HOME' ENTERED AT 17:39:35 ON 06 SEP 2006

=> File Medline EMBASE Biosis Caplus

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	0.21	0.21

FILE 'MEDLINE' ENTERED AT 17:40:06 ON 06 SEP 2006

FILE 'EMBASE' ENTERED AT 17:40:06 ON 06 SEP 2006

Copyright (c) 2006 Elsevier B.V. All rights reserved.

FILE 'BIOSIS' ENTERED AT 17:40:06 ON 06 SEP 2006

Copyright (c) 2006 The Thomson Corporation

FILE 'CAPLUS' ENTERED AT 17:40:06 ON 06 SEP 2006

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2006 AMERICAN CHEMICAL SOCIETY (ACS)

=> S ((c or campylobacter) (W) jejuni) (6a) sialyltransferase

L1 19 ((C OR CAMPYLOBACTER) (W) JEJUNI) (6A)

SIALYLTRANSFERASE

=> duplicate

ENTER REMOVE, IDENTIFY, ONLY, OR (?):remove

ENTER L# LIST OR (END):11

DUPLICATE PREFERENCE IS 'MEDLINE, EMBASE, BIOSIS, CAPLUS'

KEEP DUPLICATES FROM MORE THAN ONE FILE? Y/(N):n

PROCESSING COMPLETED FOR L1

L2 14 DUPLICATE REMOVE L1 (5 DUPLICATES REMOVED)

=> d 12 1-14 bib ab

L2 ANSWER 1 OF 14 CAPLUS COPYRIGHT 2006 ACS on STN

AN 2006:272843 CAPLUS

DN 144:326938

TI Conserved protein sequence motifs for bacterial
sialyltransferases and
uses thereof

IN Gilbert, Michel; Wakarchuk, Warren W.

PA National Research Council of Canada, Can.

SO PCT Int. Appl., 88 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.
DATE			
-----	----	-----	-----
PI WO 2006029538	A1	20060323	WO 2005-CA1432
20050916			
WO 2006029538	C1	20060601	
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ,			
CA, CH,			
CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,			
GB, GD,			
GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP,			
KR, KZ,			
LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW,			
MX, MZ,			
NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD,			
SE, SG,			
SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,			
VC, VN,			
YU, ZA, ZM, ZW			
RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR,			
HU, IE,			
IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR,			
BF, BJ,			
CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG,			
BW, GH,			
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM,			
AZ, BY,			
KG, KZ, MD, RU, TJ, TM			
PRAI US 2004-610807P	P	20040917	
AB The invention provides sialyltransferases comprising conserved protein sequence motifs, from Campylobacter jejuni strains O:36 and O:19 and Haemophilus influenzae. The sialyltransferases include α -2,3-sialyltransferase and α -2,8-sialyltransferase activities. The invention also claims methods of making sialylated products, including oligosaccharides, glycolipids, glycopeptides, or glycoproteins, using those sialyltransferases. Campylobacter jejuni CstI			

enzymes were expressed in Escherichia coli and assayed for α 2,3-sialyltransferase activity using CMP-Neu5Ac as the donor and

Lac-FCHASE (6-(5-fluorescein-carboxamido)-hexanoic acid succimidyl ester) as acceptor.

RE.CNT 8 THERE ARE 8 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L2 ANSWER 2 OF 14 CAPLUS COPYRIGHT 2006 ACS on STN

AN 2006:503091 CAPLUS

DN 145:209308

TI Bacterial sialyltransferases for carbohydrate synthesis

AU Schwardt, Oliver; Visekruna, Tamara; Rabbani, Said; Ernst, Beat

CS Institute of Molecular Pharmacy, University of Basel, Basel, CH-4056,

Switz.

SO Chimia (2006), 60(4), 234-240

CODEN: CHIMAD; ISSN: 0009-4293

PB Swiss Chemical Society

DT Journal; General Review

LA English

AB A review. Sialylation catalyzed by sialyltransferases is one of the most

interesting enzymic glycosyl transfer reactions, since chemical sialylations

usually give only low yields and lead to poor stereoselectivities. In the

last decade, several bacterial sialyltransferases were identified and

found to exhibit broader substrate specificity than their mammalian

counterparts. This suggests the potential usefulness of bacterial

sialyltransferases in chemo-enzymic synthesis of natural and non-natural

sialooligosaccharides.

RE.CNT 46 THERE ARE 46 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L2 ANSWER 3 OF 14 CAPLUS COPYRIGHT 2006 ACS on STN

AN 2005:497944 CAPLUS

DN 144:101691

TI Genomic diversity in Campylobacter jejuni: identification of C. jejuni

81-176-specific genes

AU Poly, Frederic; Threadgill, Deborah; Stintzi, Alain

CS Department of Veterinary Pathobiology, College of Veterinary Medicine,

Oklahoma State University, Stillwater, OK, 74078, USA

SO Journal of Clinical Microbiology (2005), 43(5), 2330-2338

CODEN: JCMIDW; ISSN: 0095-1137

PB American Society for Microbiology

DT Journal

LA English

AB Since the publication of the complete genomic sequence of *Campylobacter*

jejuni NCTC 11168 in Feb. 2000, evidence has been compiling that suggests

C. jejuni strains exhibit high genomic diversity. In order to investigate

this diversity, the unique genomic DNA sequences from a nonsequenced

Campylobacter strain, *C. jejuni* 81-176, were identified by comparison with

C. jejuni NCTC 11168 by using a shotgun DNA microarray approach.

Up to 63

kb of new chromosomal DNA sequences unique to this pathogen were obtained.

Eighty-six open reading frames were identified by the presence of uninterrupted coding regions encoding a min. of 40 amino acids.

In addition,

this study shows that the whole-plasmid shotgun microarray approach is

effective and provides a comprehensive coverage of DNA regions that differ

between two closely related genomes. The two plasmids harbored by this

Campylobacter strain, pTet and pVir, were also sequenced, with coverages

of 2.5- and 2.9-fold, resp., representing 72 and 92% of their complete

nucleotide sequences. The unique chromosomal genes encode proteins

involved in capsule and lipooligosaccharide biosynthesis, restriction and

modification systems, and respiratory metabolism. Several of these unique

genes are likely associated with *C. jejuni* 81-176 fitness and virulence.

Interestingly, the comparison of *C. jejuni* 81-176 unique genes with those

of *C. jejuni* ATCC 43431 revealed a single gene which encodes a probable

TraG-like protein. The product of this gene might be associated with the

mechanism of *C. jejuni* invasion into epithelial cells. In conclusion,

this study extends the repertoire of *C. jejuni* genes and thus will permit

the construction of a composite and more comprehensive microarray of *C.*

jejuni.

RE.CNT 38 THERE ARE 38 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L2 ANSWER 4 OF 14 CAPLUS COPYRIGHT 2006 ACS on STN

AN 2005:200036 CAPLUS

DN 142:428819

TI Highly efficient biosynthesis of the oligosaccharide moiety of
the GD3

ganglioside by using metabolically engineered Escherichia coli

AU Antoine, Tatiana; Heyraud, Alain; Bosso, Claude; Samain, Eric

CS CERMAV-CNRS, Grenoble, 38041, Fr.

SO Angewandte Chemie, International Edition (2005), 44(9),
1350-1352,

S1350/1-S1350/5

CODEN: ACIEF5; ISSN: 1433-7851

PB Wiley-VCH Verlag GmbH & Co. KGaA

DT Journal

LA English

AB Express order for oligosaccharides: A microbiol. process for the
synthesis

of the carbohydrate portion of gangliosides GD3 and GT3 is
described.

Lactose and sialic acid are used as exogenous precursors by a
metabolically engineered Escherichia coli strain that
overexpresses the

bifunctional sialyltransferase cstII gene from
Campylobacter jejuni.

RE.CNT 15 THERE ARE 15 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L2 ANSWER 5 OF 14 BIOSIS COPYRIGHT (c) 2006 The Thomson
Corporation on STN

AN 2004:151684 BIOSIS

DN PREV200400154694

TI Lipopolysaccharide alpha-2,3 sialyltransferase of
Campylobacter jejuni and its uses.

AU Gilbert, Michel [Inventor, Reprint Author]; Wakarchuk, Warren W.
[Inventor]

CS Hull, Canada

ASSIGNEE: National Research Council of Canada, Ottawa, Canada

PI US 6689604 20040210

SO Official Gazette of the United States Patent and Trademark
Office Patents,

(Feb 10 2004) Vol. 1279, No. 2.

<http://www.uspto.gov/web/menu/patdata.html>

. e-file.

ISSN: 0098-1133 (ISSN print).

DT Patent

LA English

ED Entered STN: 17 Mar 2004

Last Updated on STN: 17 Mar 2004

AB The structure and specificity of a recombinant
alpha2,3-sialyltransferase
from *Campylobacter* spp., is disclosed. Also provided are
methods for
using the alpha2,3-sialyltransferase in the production of desired
carbohydrate structures and nucleic acids that encode the
sialyltransferase.

L2 ANSWER 6 OF 14 BIOSIS COPYRIGHT (c) 2006 The Thomson
Corporation on STN
AN 2005:321982 BIOSIS
DN PREV200510111764
TI Domain organization of the Cst-I sialyltransferase from
Campylobacter jejuni.
AU Gilbert, Michel [Reprint Author]; Karwaski, Marie-France;
Brochu, Denis;
Wakarchuk, Warren W.
CS Natl Res Council Canada, Inst Biol Sci, Ottawa, ON K1A 0R6,
Canada
SO Glycobiology, (NOV 2004) Vol. 14, No. 11, pp. 1126.
Meeting Info.: Joint Meeting of the
Society-for-Glycobiology/Japanese-
Society-for-Carbohydrate-Research. Honolulu, HI, USA. November
17 -20,
2004. Soc Gylcobiol; Japanese Soc Carbohydrate Res.
ISSN: 0959-6658.
DT Conference; (Meeting)
Conference; Abstract; (Meeting Abstract)
LA English
ED Entered STN: 25 Aug 2005
Last Updated on STN: 25 Aug 2005

L2 ANSWER 7 OF 14 BIOSIS COPYRIGHT (c) 2006 The Thomson
Corporation on STN
AN 2005:321973 BIOSIS
DN PREV200510111755
TI Towards the understanding of the catalytic mechanism and
substrate
specificities of sialyltransferases from *Campylobacter*
jejuni.
AU Chiu, Cecilia P. C. [Reprint Author]; Gilbert, Michel; Lairson,
Luke L.;
Watts, Andrew; Wakarchuk, Warren W.; Withers, Stephen G.;
Strynadka,
Natalie C. J.
CS Univ British Columbia, Dept Biochem and Mol Biol, Vancouver, BC
V6T 1Z3,
Canada
SO Glycobiology, (NOV 2004) Vol. 14, No. 11, pp. 1123.
Meeting Info.: Joint Meeting of the
Society-for-Glycobiology/Japanese-

Society-for-Carbohydrate-Research. Honolulu, HI, USA. November
17 -20,

2004. Soc Glycobiol; Japanese Soc Carbohydrate Res.
ISSN: 0959-6658.

DT Conference; (Meeting)
Conference; Abstract; (Meeting Abstract)

LA English

ED Entered STN: 25 Aug 2005

Last Updated on STN: 25 Aug 2005

L2 ANSWER 8 OF 14 MEDLINE on STN DUPLICATE 1

AN 2004048991 MEDLINE

DN PubMed ID: 14730352

TI Structural analysis of the sialyltransferase CstII from
Campylobacter jejuni in complex with a substrate analog.

AU Chiu Cecilia P C; Watts Andrew G; Lairson Luke L; Gilbert
Michel; Lim

Daniel; Wakarchuk Warren W; Withers Stephen G; Strynadka Natalie
C J

CS Department of Biochemistry and Molecular Biology, University of
British

Columbia, 2146 Health Sciences Mall, Vancouver, British Columbia

V6T 1Z3,

Canada.

SO Nature structural & molecular biology, (2004 Feb) Vol. 11, No.
2, pp.

163-70. Electronic Publication: 2004-01-18.

Journal code: 101186374. ISSN: 1545-9993.

CY United States

DT Journal; Article; (JOURNAL ARTICLE)

LA English

FS Priority Journals

OS PDB-1RO7; PDB-1RO8

EM 200404

ED Entered STN: 30 Jan 2004

Last Updated on STN: 6 Apr 2004

Entered Medline: 5 Apr 2004

AB Sialic acid terminates oligosaccharide chains on mammalian and
microbial

cell surfaces, playing critical roles in recognition and
adherence. The

enzymes that transfer the sialic acid moiety from
cytidine-5'-monophospho-

N-acetyl-neuraminic acid (CMP-NeuAc) to the terminal positions
of these

key glycoconjugates are known as sialyltransferases. Despite
their

important biological roles, little is understood about the
mechanism or

molecular structure of these membrane-associated enzymes. We
report the

first structure of a sialyltransferase, that of CstII from
 Campylobacter jejuni, a highly prevalent foodborne
 pathogen. Our structural, mutagenesis and kinetic data provide
 support
 for a novel mode of substrate binding and glycosyl transfer
 mechanism,
 including essential roles of a histidine (general base) and two
 tyrosine
 residues (coordination of the phosphate leaving group). This
 work
 provides a framework for understanding the activity of several
 sialyltransferases, from bacterial to human, and for the
 structure-based
 design of specific inhibitors.

L2 ANSWER 9 OF 14 CAPLUS COPYRIGHT 2006 ACS on STN

AN 2002:276514 CAPLUS

DN 136:320378

TI Campylobacter glycosyltransferase genes and enzymes for
 biosynthesis of

gangliosides and ganglioside mimics

IN Gilbert, Michel; Wakarchuk, Warren W.

PA National Research Council of Canada, Can.

SO U.S. Pat. Appl. Publ., 84 pp., Cont.-in-part of U.S. Ser. No.
 495,406.

CODEN: USXXCO

DT Patent

LA English

FAN.CNT 3

	PATENT NO.	KIND	DATE	APPLICATION NO.
DATE	-----	----	-----	-----
PI	US 2002042369	A1	20020411	US 2001-816028
20010321				
	US 6699705	B2	20040302	
	US 6503744	B1	20030107	US 2000-495406
20000131				
	EP 1652927	A2	20060503	EP 2005-25316
20000201				
	EP 1652927	A3	20060719	
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE,			
MC, PT,				
	IE, SI, LT, LV, FI, RO, MK, CY, AL			
	AT 329036	E	20060615	AT 2000-901455
20000201				
	CA 2441570	AA	20020926	CA 2002-2441570
20020222				
	WO 2002074942	A2	20020926	WO 2002-CA229
20020222				
	WO 2002074942	A3	20030313	

WO 2002074942	B1	20030703	
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA,			
CH, CN,			
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD,			
GE, GH,			
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,			
LK, LR,			
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ,			
OM, PH,			
PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR,			
TT, TZ,			
UA, UG, US, UZ, VN, YU, ZA, ZM, ZW			
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM,			
AZ, BY,			
KG, KZ, MD, RU, TJ, TM, AT, BE, CH, CY, DE, DK, ES, FI,			
FR, GB,			
GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI,			
CM, GA,			
GN, GQ, GW, ML, MR, NE, SN, TD, TG			
EP 1385941	A2	20040204	EP 2002-703414
20020222			
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE,			
MC, PT,			
IE, SI, LT, LV, FI, RO, MK, CY, AL, TR			
JP 2004524033	T2	20040812	JP 2002-574334
20020222			
US 2003148459	A1	20030807	US 2002-303161
20021121			
US 2003157655	A1	20030821	US 2002-303118
20021121			
US 6905867	B2	20050614	
US 2003157656	A1	20030821	US 2002-303128
20021121			
US 6911337	B2	20050628	
US 2003157657	A1	20030821	US 2002-303134
20021121			
US 6825019	B2	20041130	
US 2003157658	A1	20030821	US 2002-303162
20021121			
US 6723545	B2	20040420	
US 2004180406	A1	20040916	US 2003-735419
20031211			
US 7026147	B2	20060411	
US 2006166317	A1	20060727	US 2003-734719
20031211			
US 2004203103	A1	20041014	US 2004-820536
20040407			
US 2004229313	A1	20041118	US 2004-821573
20040408			
US 2004229263	A1	20041118	US 2004-821604
20040408			

US 2004265875	A1	20041230	US 2004-830825
20040424			
US 2004203112	A1	20041014	US 2004-845408
20040512			
US 2004203113	A1	20041014	US 2004-845412
20040512			
US 2004219638	A1	20041104	US 2004-846219
20040514			
US 2004229272	A1	20041118	US 2004-847983
20040517			
US 2004259203	A1	20041223	US 2004-850125
20040519			
US 2004259140	A1	20041223	US 2004-850807
20040521			
US 2005048630	A1	20050303	US 2004-962334
20041008			
US 2005084891	A1	20050421	US 2004-962235
20041008			
US 2005227248	A1	20051013	US 2004-961882
20041008			
US 7078207	B2	20060718	
PRAI US 1999-118213P	P	19990201	
US 2000-495406	A2	20000131	
EP 2000-901455	A3	20000201	
US 2001-816028	A	20010321	
WO 2002-CA229	W	20020222	
US 2002-303118	A3	20021121	
US 2002-303128	A1	20021121	
US 2002-303134	A3	20021121	

AB This invention provides *Campylobacter jejuni* glycosyltransferases, including a bifunctional sialyltransferase that has both an α 2,3- and an α 2,8-activity. A β 1,4-GaINAc transferase and a β 1,3-galactosyltransferase are also provided by the invention, as are other glycosyltransferases and enzymes involved in synthesis of lipooligosaccharide (LOS). In addnl.

embodiments, the invention provides nucleic acids that encode the glycosyltransferases, as well as expression vectors and host cells for expressing the glycosyltransferases. The enzymes may be used in preparation of gangliosides, lysogangliosides, and mimics of gangliosides and lysogangliosides. Thus, *C. jejuni* gene cstI α 2,3- sialyltransferase, gene cstII bifunctional α 2,3/ α 2,8-sialyltransferase, gene cgtA β -1,4-N-acetylgalactosaminyltransferase, and gene cgtB β -1,3-galactosyltransferase enzymes were used to prepare the carbohydrate portion of gangliosides GM1a, GM2, GM3, GD1a, GD3, and GT1a.

L2 ANSWER 10 OF 14 CAPLUS COPYRIGHT 2006 ACS on STN

AN 2001:435090 CAPLUS

DN 135:5768

TI Synthesis of sialylated oligosaccharide donors via sialylation
and enzymic
glycosidation

IN Mehta, Seema; Gilbert, Michel; Wakarchuk, Warren W.; Whitfield,
Dennis M.

PA National Research Council of Canada, Can.

SO PCT Int. Appl., 35 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.
DATE	-----	----	-----	-----

PI WO 2001042264 A1 20010614 WO 2000-CA1487

20001208

W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA,
CH, CN,
CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH,
GM, HR,
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT,
LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT,
RO, RU,
SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US,
UZ, VN,
YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE,
CH, CY,
DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE,
TR, BF,
BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG

PRAI US 1999-169945P P 19991210

AB A method for the synthesis of aryl thio glycosides comprising a
sialylated

residue of β -D-galactose is disclosed. The method consists of
preparing

by a chemical synthesis a non-sialylated aryl thio glycoside,
and enzymically

sialylating the latter with a sialic acid in the presence of a
suitable

sialyltransferase. The sialylated aryl thio glycoside is then
chemical

derivatized by standard procedures, to provide a derivative
suitable for use as a

donor in chemical syntheses of sialylated oligosaccharides. The
derivatized

sialylated aryl thio glycosides are prepared in high yields, due to reduced number of chemical and purification steps involved in the process. Derivatized aryl thio glycosides useful as building blocks for the synthesis of biol. active sialylated oligosaccharides are also disclosed. Thus, [Methyl (5-acetamido-4,7,8,9-tetra-O-acetyl-3,5-dideoxy-D-glycero- α -D-galacto-2-nonulopyranosyl)onate]-(2,3)-O-(2,4,6-tri-O-acetyl- β -D-galactopyranosyl)-(1,4)-3-O-acetyl-6-O-tert-butyl-diphenylsilyl-2-deoxy-2-phthalimido- β -D-glucopyranoside was prepared via sialylation and enzymic glycosidation reactions.

RE.CNT 7 THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L2 ANSWER 11 OF 14 CAPLUS COPYRIGHT 2006 ACS on STN
AN 2000:553711 CAPLUS
DN 133:161277
TI Campylobacter glycosyltransferases for biosynthesis of gangliosides and ganglioside mimics
IN Gilbert, Michel; Wakarchuk, Warren W.
PA National Research Council of Canada, Can.
SO PCT Int. Appl., 120 pp.
CODEN: PIXXD2
DT Patent
LA English
FAN.CNT 3

PATENT NO.	KIND	DATE	APPLICATION NO.
WO 2000046379	A1	20000810	WO 2000-CA86
20000201			
W:	AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, ZA, TJ, TM		
RW:	GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF,		

CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG
 US 6503744 B1 20030107 US 2000-495406
 20000131
 CA 2360205 AA 20000810 CA 2000-2360205
 20000201
 EP 1147200 A1 20011024 EP 2000-901455
 20000201
 EP 1147200 B1 20060607
 R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE,
 MC, PT,
 IE, LT, LV, FI, RO, CY
 JP 2002535992 T2 20021029 JP 2000-597438
 20000201
 AU 772569 B2 20040429 AU 2000-22743
 20000201
 EP 1652927 A2 20060503 EP 2005-25316
 20000201
 EP 1652927 A3 20060719
 R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE,
 MC, PT,

IE, SI, LT, LV, FI, RO, MK, CY, AL
 AT 329036 E 20060615 AT 2000-901455
 20000201
 AU 2004203474 A1 20040826 AU 2004-203474
 20040729
 PRAI US 1999-118213P P 19990201
 US 2000-495406 A 20000131
 EP 2000-901455 A3 20000201
 WO 2000-CA86 W 20000201

AB This invention provides prokaryotic glycosyltransferases,
 including a
 bifunctional sialyltransferase that has both an α 2,3- and an
 α 2,8- activity. A β 1,4-GalNAc transferase and a
 β 1,3-galactosyltransferase are also provided by the invention,
 as are
 other glycosyltransferases and enzymes involved in synthesis of
 lipooligosaccharide (LOS). The glycosyltransferases can be
 obtained from,
 for example, Campylobacter species, including C. jejuni. In
 addnl.
 embodiments, the invention provides nucleic acids that encode the
 glycosyltransferases, as well as expression vectors and host
 cells for
 expressing the glycosyltransferases.

RE.CNT 11 THERE ARE 11 CITED REFERENCES AVAILABLE FOR THIS RECORD
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L2 ANSWER 12 OF 14 BIOSIS COPYRIGHT (c) 2006 The Thomson
 Corporation on
 STN
 AN 2001:93201 BIOSIS

DN PREV200100093201
 TI Modulation of the mono- and bi-functional activity of the
 Campylobacter jejuni Cst-II sialyltransferase:
 A novel phase variation mechanism.
 AU Gilbert, Michel [Reprint author]; Karwaski, Marie-France
 [Reprint author];
 Cunningham, Anna-Maria [Reprint author]; Wakarchuk, Warren W.
 [Reprint
 author]
 CS Institute for Biological Sciences, NRCC, 100 Sussex Dr., Ottawa,
 ON, K1A
 0R6, Canada
 SO Glycoconjugate Journal, (January-February, 2000) Vol. 17, No.
 1-2, pp. 91.
 print.
 Meeting Info.: Second International Glycosyltransferase
 Symposium.
 Toronto, Ontario, Canada. May 12-14, 2000.
 ISSN: 0282-0080.
 DT Conference; (Meeting)
 Conference; Abstract; (Meeting Abstract)
 LA English
 ED Entered STN: 21 Feb 2001
 Last Updated on STN: 12 Feb 2002

L2 ANSWER 13 OF 14 CAPLUS COPYRIGHT 2006 ACS on STN
 AN 1999:626342 CAPLUS
 DN 131:253359
 TI Campylobacter jejuni gene cst-I lipopolysaccharide α -2,3
 sialyltransferase, its DNA and amino acid sequences, recombinant
 production, and its acceptor specificity
 IN Gilbert, Michel; Wakarchuk, Warren W.
 PA National Research Council of Canada, Can.
 SO PCT Int. Appl., 47 pp.
 CODEN: PIXXD2
 DT Patent
 LA English
 FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.
DATE			
-----	----	-----	-----

PI WO 9949051	A1	19990930	WO 1999-CÅ238
19990322			
W: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN,			
CU, CZ,			
DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL,			
IN, IS,			
JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD,			
MG, MK,			
MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK,			
SL, TJ,			

TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG,
KZ, MD,

RU, TJ, TM
RW: GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW, AT, BE, CH, CY,
DE, DK,

ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ,
CF, CG,

CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG
US 6689604 B1 20040210 US 1999-272960

19990318
CA 2323753 AA 19990930 CA 1999-2323753

19990322
AU 9928230 A1 19991018 AU 1999-28230

19990322
AU 745040 B2 20020307

EP 1082440 A1 20010314 EP 1999-908717
19990322

R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE,
MC, PT,

IE, FI
JP 2002507424 T2 20020312 JP 2000-538012
19990322

US 2003049270 A1 20030313 US 2002-58636
20020129

US 6709834 B2 20040323

US 2004152165 A1 20040805 US 2004-799016
20040311

PRAI US 1998-78891P P 19980320

US 1999-272960 A 19990318

WO 1999-CA238 W 19990322

US 2002-58636 A3 20020129
AB The invention provides DNA mols. that encode gene cst-I
lipopolysaccharide

α -2,3 sialyltransferase of Campylobacter
jejuni. The DNA sequence of C. jejuni gene cst-I, as well as the
corresponding amino acid sequence of lipopolysaccharide α -2,3
sialyltransferase are claimed. The invention also provides

methods for
the recombinant production of lipopolysaccharide α -2,3
sialyltransferase

in prokaryotic and eukaryotic cells. The invention further
provides the

specificity of the C. jejuni lipopolysaccharide
 α -2,3 sialyltransferase. The C. jejuni

lipopolysaccharide α -2,3 sialyltransferase uses terminal
galactose acceptors that are β -(1 \rightarrow 4) linked to either glucose
or N-acetylglucosamine. The enzyme also uses terminal galactose

acceptors
that are β -(1 \rightarrow 3) linked to N-acetylglucosamine or
N-acetylgalactosamine. The enzyme uses cytidine monophosphate-N-

acetylneuraminic acid (CMP-Neu5Ac) as the donor. The broad
acceptor

specificity of lipopolysaccharide α -2,3 sialyltransferase encoded by cst-I demonstrates its utility and makes it an attractive tool for chemo-enzymic synthesis of sialylated oligosaccharides.

RE.CNT 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L2 ANSWER 14 OF 14 MEDLINE on STN DUPLICATE 2
AN 1999449955 MEDLINE
DN PubMed ID: 10520252
TI Synthesis of a disialylated hexasaccharide of type VIII group B Streptococcus capsular polysaccharide.
AU Eichler E; Jennings H J; Gilbert M; Whitfield D M
CS National Research Council, Ottawa, Ontario, Canada.
SO Carbohydrate research, (1999 Jun 30) Vol. 319, No. 1-4, pp. 1-16.
 Journal code: 0043535. ISSN: 0008-6215.
CY Netherlands
DT Journal; Article; (JOURNAL ARTICLE)
LA English
FS Priority Journals
EM 199912
ED Entered STN: 13 Jan 2000
 Last Updated on STN: 13 Jan 2000
 Entered Medline: 17 Dec 1999
AB As part of our program to design, develop and prepare protective vaccines against the bacterial pathogens Group B Streptococcus, we report the synthesis of a disialylated hexasaccharide. This hexasaccharide represents a portion of the serotype-specific capsular polysaccharide of Type VIII that has the tetrasaccharide repeat unit [beta-L-Rhap-(1-->4)-beta-D-Glcp-(1-->4)-[alpha-Neu5Ac-(2--> 3)]-beta-D-Galp-(1-->4)]n. A tetrasaccharide corresponding to this repeat unit has been synthesized by us [E. Eichler, H.J. Jennings, D.M. Whitfield, J. Carbohydr. Chemical, 16 (1997) 385-411]. Since the protective epitopes are believed to involve several repeat units, methods to extend this tetrasaccharide were examined. This objective requires a glycosylation of the unreactive OH-4 of the beta-L-Rhap, which was accomplished by coupling a D-Galp glycosyl trichloroacetimidate donor with a beta-L-Rhap-(1-->4)-D-Glcp acceptor. Subsequent coupling of this trisaccharide as a donor to an

alpha-Neu5Ac-(2-->3)-D-Galp disaccharide acceptor gave a pentasaccharide.

The pentasaccharide was deprotected and enzymatically sialylated using an

alpha-(2-->3)-sialyltransferase from *Campylobacter*

jejuni to give the title hexasaccharide alpha-Neu5Ac-(2-->3)-

beta-D-Galp-(1-->4)-beta-L-Rhap-(1-->4)-beta-D-Glcp-(1-->4)-[alpha-Neu5Ac-(2-->3)]-beta-D-Galp-(1-->O)-(CH₂)₃N₃.